STD PREVENTION AND CONTROL SERVICES, SAN FRANCISCO DEPARTMENT OF PUBLIC HEALTH STD AND HIV SCREENING GUIDELINES

These evidence-based recommendations provide guidance for chlamydia, gonorrhea, syphilis and HIV screening in persons without symptoms or a need for diagnostic testing.¹

WOMEN

| | Chlamydia and Gonorrhea | Syphilis | HIV |
|---|--|---|---|
| 25 years and younger | Test every 12 months | Not routinely recommended ² | At least one lifetime test ² |
| Older than 25 years | Not routinely recommended ² | | At least one lifetime test ² |
| Pregnant ³ | Test in 1 st trimester, repeat in 3 rd trimester if at increased risk ² | Test in 1 st and 3 rd trimester, and at delivery ⁴ | First prenatal visit, repeat in 3 rd trimester if at increased risk ² |
| HIV+ Women should be screened annually for trichomoniasis (trich) | | | |

MEN WHO HAVE SEX WITH WOMEN

| | Chlamydia and Gonorrhea | Syphilis | HIV |
|-------------------|--|----------|---|
| Any age, any site | Not routinely recommended ² | | At least one lifetime test ² |

MEN WHO HAVE SEX WITH MEN (MSM)

| ment with the sex with ment (mon) | | | | | |
|-----------------------------------|-------------------------|----------------|-----|--|--|
| | Chlamydia and Gonorrhea | Syphilis | HIV | | |
| Blood | | Every 3 months | | | |
| Rectal & Pharyngeal ⁵ | Every 3 months | | | | |
| Urine ⁶ | Every 3 months | | | | |

TRANS WOMEN AND TRANS MEN WHO HAVE SEX WITH MEN

| | Chlamydia and Gonorrhea | Syphilis | HIV |
|----------------------------------|-------------------------|----------------|-----|
| Blood | | Every 3 months | |
| Rectal & Pharyngeal ⁵ | Every 3 months | | |
| Urine ⁶ /Vaginal Swab | Every 3 months | | |

¹These are general population recommendations. SFDPH may make separate recommendations for specific groups that correlate with higher risk.

Terminology Defined: Transgender (Trans): A person whose gender identity differs from the sex that was assigned at birth. A trans man (or trans male) is someone with a male gender identity who was assigned female sex at birth; a trans woman (or trans female) is someone with a female gender identity who was assigned male sex at birth.

For more info: Visit www.sfcityclinic.org or call 415-487-5595

²Screen if patient has any of the following risk factors: sex with a man who has sex with men, history of STD in the past year, methamphetamine use, unstable housing or homelessness, sex work, intimate partner violence, or incarceration.

³Regardless of intentions to carry to term.

⁴CDPH Expanded Syphilis Screening Recommendations for the Prevention of Congenital Syphilis: 2020

⁵Prioritize these extra-genital sites, as rectal and pharyngeal infections are almost always asymptomatic.

⁶If cost permits or extra-genital testing not available.